































Poor concentration, seeing things	Poor concentration, seeing things	Poor concentration, seeing things	Poor concentration, seeing things
Agitation	Agitation	Agitation	Agitation
Talking to themselves, confusion	Talking to themselves, confusion	Talking to themselves, confusion	Talking to themselves, confusion
Hypoactive (drowsy, sleepy)	Hypoactive (drowsy, sleepy)	Hypoactive (drowsy, sleepy)	Hypoactive (drowsy, sleepy)
Hyperactive (restless, agitated)	Hyperactive (restless, agitated)	Hyperactive (restless, agitated)	Hyperactive (restless, agitated)

| Poor concentration, seeing things |
|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|
| Agitation | Agitation | Agitation | Agitation |
| Talking to themselves, confusion |
| Hypoactive (drowsy, sleepy) | Hypoactive (drowsy, sleepy) | Hypoactive (drowsy, sleepy) | Hypoactive (drowsy, sleepy) |
| Hyperactive
(restless, agitated) | Hyperactive
(restless, agitated) | Hyperactive
(restless, agitated) | Hyperactive
(restless, agitated) |

_ _

































| Distress, feeling of impending dread |
|---|---|---|---|
| Rapid breathing | Rapid breathing | Rapid breathing | Rapid breathing |
| Agitation | Agitation | Agitation | Agitation |
| Sweating, nausea, vomiting | Sweating, nausea, vomiting | Sweating, nausea, vomiting | Sweating, nausea, vomiting |
| Heavy, tight,
crushing feeling in
chest |
| Indigestion-type pain, pain in jaw/arm |

| Distress, feeling of impending dread |
|---|---|---|---|
| Rapid breathing | Rapid breathing | Rapid breathing | Rapid breathing |
| Agitation | Agitation | Agitation | Agitation |
| Sweating, nausea, vomiting | Sweating, nausea, vomiting | Sweating, nausea, vomiting | Sweating, nausea, vomiting |
| Heavy, tight,
crushing feeling in
chest |
| Indigestion-type pain, pain in jaw/arm |



(Difficulty Breathing)



Dyspnoea





| Breathing quickly or noisily |
|--|--|--|--|
| Agitation, confusion | Agitation, confusion | Agitation, confusion | Agitation, confusion |
| Sweating, fever | Sweating, fever | Sweating, fever | Sweating, fever |
| Short of breath at rest or on movement | Short of breath at rest or on movement | Short of breath at rest or on movement | Short of breath at rest or on movement |
| Cough (dry or moist) |

| Breathing quickly or noisily |
|--|--|--|--|
| Agitation, confusion | Agitation, confusion | Agitation, confusion | Agitation, confusion |
| Sweating, fever | Sweating, fever | Sweating, fever | Sweating, fever |
| Short of breath at rest or on movement | Short of breath at rest or on movement | Short of breath at rest or on movement | Short of breath at rest or on movement |
| Cough (dry or moist) |

































| Dry tongue, lips and skin |
|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|
| Constipation, diarrhea, vomiting | Constipation, diarrhea, vomiting | Constipation, diarrhea, vomiting | Constipation, diarrhea, vomiting |
| Decreased bodily fluids, dark urine |
| Agitation/confusion | Agitation/confusion | Agitation/confusion | Agitation/confusion |
| Fatigue, unwilling to do activities |

| Dry tongue, lips and skin |
|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|
| Constipation, diarrhea, vomiting | Constipation, diarrhea, vomiting | Constipation, diarrhea, vomiting | Constipation, diarrhea, vomiting |
| Decreased bodily fluids, dark urine |
| Agitation/confusion | Agitation/confusion | Agitation/confusion | Agitation/confusion |
| Fatigue, unwilling to do activities |

























Agitation, restlessness, confusion	Agitation, restlessness, confusion	Agitation, restlessness, confusion	Agitation, restlessness, confusion
Pain/burning when passing urine			
Fatigue, tiredness, lack of appetite			
Smelly, cloudy milky or bloody urine			
Increased toileting, incontinence aid or bed changes, can't control or pass urine	Increased toileting, incontinence aid or bed changes, can't control or pass urine	Increased toileting, incontinence aid or bed changes, can't control or pass urine	Increased toileting, incontinence aid or bed changes, can't control or pass urine

Agitation, restlessness, confusion	Agitation, restlessness, confusion	Agitation, restlessness, confusion	Agitation, restlessness, confusion
Pain/burning when passing urine			
Fatigue, tiredness, lack of appetite			
Smelly, cloudy milky or bloody urine			
Increased toileting, incontinence aid or bed changes, can't control or pass urine	Increased toileting, incontinence aid or bed changes, can't control or pass urine	Increased toileting, incontinence aid or bed changes, can't control or pass urine	Increased toileting, incontinence aid or bed changes, can't control or pass urine


































Stools less than 3 times a week, more than 2 days without passing stool	Stools less than 3 times a week, more than 2 days without passing stool	Stools less than 3 times a week, more than 2 days without passing stool Stools less than 3 times a week, more than 2 days without passing stool
Hard and dry pellets, stool stuck or unable to pass	Hard and dry pellets, stool stuck or unable to pass	Hard and dry pellets, stool stuck or unable to pass Hard and dry pellets, stool stuck or unable to pass
Nausea or lack or appetite	Nausea or lack or appetite	Nausea or lack or appetite Nausea or lack or
Feeling of fullness or cramps	Feeling of fullness or cramps	Feeling of fullness or cramps Feeling of fullness or cramps
Agitation, restlessness	Agitation, restlessness	Agitation, Agitation, restlessness restlessness

Stools less than 3 times a week, more than 2 days without passing stool	Stools less than 3 times a week, more than 2 days without passing stool	Stools less than 3 times a week, more than 2 days without passing stool Stools less than 3 times a week, more than 2 days without passing stool
Hard and dry pellets, stool stuck or unable to pass	Hard and dry pellets, stool stuck or unable to pass	Hard and dry pellets, stool stuck or unable to pass Hard and dry pellets, stool stuck or unable to pass
Nausea or lack or appetite	Nausea or lack or appetite	Nausea or lack or appetite Nausea or lack or
Feeling of fullness or cramps	Feeling of fullness or cramps	Feeling of fullness or cramps Feeling of fullness or cramps
Agitation, restlessness	Agitation, restlessness	Agitation, Agitation, restlessness restlessness




































Recent illness, frailty, delirium, or return from hospital	Recent illness, frailty, delirium, or return from hospital
New to the home	New to the home
Decreased walking and activity recently	Decreased walking and activity recently
Resident not wearing glasses, hearing aids or footwear	Resident not wearing glasses, hearing aids or footwear
Not using mobility aids	Not using mobility aids

Recent illness, frailty, delirium, or return from hospital	Recent illness, frailty, delirium, or return from hospital	
New to the home	New to the home	
Decreased walking and activity recently	Decreased walking and activity recently	
Resident not wearing glasses, hearing aids or footwear	Resident not wearing glasses, hearing aids or footwear	
Not using mobility aids	Not using mobility aids	

Recent illness, frailty, delirium, or return from hospital	Recent illness, frailty, delirium, or return from hospital
New to the home	New to the home
Decreased walking and activity recently	Decreased walking and activity recently
Resident not wearing glasses, hearing aids or footwear	Resident not wearing glasses, hearing aids or footwear
Not using mobility aids	Not using mobility aids

Recent illness, frailty, delirium, or return from hospital	Recent illness, frailty, delirium, or return from hospital	
New to the home	New to the home	
Decreased walking and activity recently	Decreased walking and activity recently	
Resident not wearing glasses, hearing aids or footwear	Resident not wearing glasses, hearing aids or footwear	
Not using mobility aids	Not using mobility aids	

































Increased agitation, confusion, tiredness	Increased agitation, confusion, tiredness	Increased agitation, confusion, tiredness	Increased agitation, confusion, tiredness
Increased pain and distress	Increased pain and distress	Increased pain and distress	Increased pain and distress
Unable to perform usual activities	Unable to perform usual activities	Unable to perform usual activities	Unable to perform usual activities
Worsening condition and disease progression	Worsening condition and disease progression	Worsening condition and disease progression	Worsening condition and disease progression
Decreasing alertness and reduced diet and fluid intake	Decreasing alertness and reduced diet and fluid intake	Decreasing alertness and reduced diet and fluid intake	Decreasing alertness and reduced diet and fluid intake

Increased agitation, confusion, tiredness	Increased agitation, confusion, tiredness	Increased agitation, confusion, tiredness	Increased agitation, confusion, tiredness
Increased pain and distress	Increased pain and distress	Increased pain and distress	Increased pain and distress
Unable to perform usual activities	Unable to perform usual activities	Unable to perform usual activities	Unable to perform usual activities
Worsening condition and disease progression	Worsening condition and disease progression	Worsening condition and disease progression	Worsening condition and disease progression
Decreasing alertness and reduced diet and fluid intake	Decreasing alertness and reduced diet and fluid intake	Decreasing alertness and reduced diet and fluid intake	Decreasing alertness and reduced diet and fluid intake